



<https://www.biodiversitylibrary.org/>

Annual report of the Ohio State Academy of Science.

Columbus :The Academy,1893-1930.

<https://www.biodiversitylibrary.org/bibliography/14675>

v.3 1901-1903: <https://www.biodiversitylibrary.org/item/50847>

Article/Chapter Title: Additions to the list of Ohio Hemiptera

Author(s): Osborn, Herbert

Subject(s): Hemiptera, Auchenorrhyncha, Fulgoroidea

Page(s): Page 36, Page 37

Holding Institution: New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by: The LuEsther T Mertz Library, the New York Botanical Garden

Generated 5 October 2018 7:54 AM

<https://www.biodiversitylibrary.org/pdf4/084084800050847>

This page intentionally left blank.

ADDITIONS TO THE LIST OF OHIO HEMIPTERA.

HERBERT OSBORN.

- Scolops dessicatus* Uh. (?). Bellaire, Aug. 28, 1900.
- Liburnia campestris* Van D. Bellaire, Aug. 28, 1900.
- Liburnia ornata* Stal. Bellaire. Common. Aug. 28, 1900.
- Clastoptera xanthocephala* Germ. Bellaire, Aug. 28, 1900. More common south.
- Idiocerus pallidus* Fh. Bellaire, Aug. 28, 1900.
- Tettigonia hartii* Wdw. (MS). Bellaire, Aug. 28, 1900.
- Gypona pectoralis* Spang. Vinton (Hine).
- Xestocephalus tessellatus* VanD. Bellaire, Aug. 28, 1900.
- Deltoccephalus obtectus* O. & B. Bellaire, Aug. 29, 1900.
- Deltoccephalus melsheimeri* Fh. Castalia.
- Deltoccephalus apicatus* Osb. Bellaire, Aug. 29, 1900. The most western point yet noted for this species, which has been found in Vermont, New Jersey, Maryland and District of Columbia.
- Athysanella acuticauda* Bak. Bellaire, Aug. 29, 1900.
- Lonatura catalina* O. & B. Bellaire, Aug. 29, 1900. The most eastern record for this species, which abounds on *Sporobulus* on the western prairies.
- Phlepsius decorus* O. & B. Bellaire, Sept. 1, 1900. Also a western species. This Bellaire example was taken on grass surrounding a small spring on a hillside.
- Phlepsius fulvidorsum* Fh. Vinton, May 5-12, 1900 (Hine).
- Thamnotettix inornata* VanD. Columbus, Sept., 1900. One example collected by Prof. E. D. Ball.
- Cicadula punctifrons* Fall. The typical european form. Bellaire, Aug. 20, 1900.
- Gnathodus abdominalis* VanD. Bellaire, Aug. 29, 1900.
- Empoasca smaragdula* Fall. Bellaire, Aug. 29-30, 1900.
- Empoasca obtusa-trifasciata* Gill.
- Dicranura flavipennis* Wooster (Webster) Bellaire.
- Typhlocyba comes-ziczzac* Walsh. Bellaire, Aug., 1900.
- Typhlocyba tricincta* Fh. Bellaire, Aug., 1900.
- Calophya nigripennis* Riley. On sumac at Sugar Grove.
- Pemphigus populi-transversus* Riley. Bellaire, Aug. 31, 1900.
- Schizoneura ulmi*. Columbus. On red elm.
- Chionaspis salicis* Harr. On willows. Bellaire, Aug. 31, 1900.
- Aspidiotus rapax* Comst. Columbus in greenhouse. (Collected by W. F. Naret.)
- Melanaethus robustus* Uhl. Bellaire, Aug. 29, 1900.
- Rhacognathus americanus* Stal. Loudonville, June 14, 1900 (Hine).

- Geocoris fuliginosus* Say. Columbus. Bellaire.
Lygaeus reclivatus Say. Sandusky. Bellaire.
Megalocoerea debilis. Uh.
Phytocoris scrupeus Say.
Compsocerocoris annulicornis. Reut. Bellaire.
Lygus invitus Say. Newark. Ashtabula.
Lygus plagiatus Uh. A western species.
Lygus monachus Uh. Columbus.
Poecyloscytus basalis Reut. Bellaire. Vinton.
Coccobaphes sanguinarius Uhl. Vinton, May 5-12, 1900 (Hine).
Mimoceps gracilis Uh. Sandusky.
Camptobrochis nebulosus Uh. Bellaire, Aug., 1900.
Eccritotarsus elegans Uh. Bellaire, Aug., 1900.
Episcopus ornatus Uh. Celina, Bellaire.
Ilnacora stalii Reut. Bellaire, Aug., 1900.
Ilnacora malina Uh. Vinton, May 1900 (Hine).
Malacocoris irroratus Say. Bellaire, Aug., 1900.
Stiprhosoma stygica Say. Bellaire, Aug., 1900.
Idolocoris agilis Uh. Vinton, May 5-12, 1900 (Hine).
Plagiognathus obscurus Uh. Bellaire, Aug., 1900.
Plagionathus sp. Bellaire, Aug., 1900.
Agaliastes associatus Uh. Bellaire. A very common and wide-spread species.
Neoborus laetus Uh. Bellaire, Aug., 1900.
Neoborus pettiti Uh. (MS.). Newark.
Aradus crenatus Say.
Sinea spinipes H. Schf.
Ectrichodia cruciata Say. Vinton. Common south.
Salda. Sandusky Bay. On a wreck about a mile from land. Aug., 1900.
Pelocoris femorata. Dredged from Black Channel, Sandusky Bay.
Corisa bivittata Prov. Dredged from shallow water, sandy bottom, off Cedar Point beach, Lake Erie.
Corisa calva Say. Columbus.